arXiv:1107.1221v4 [math.AG] 22 Jun 2012

FINITENESS OF K3 SURFACES AND THE TATE CONJECTURE

MAX LIEBLICH, DAVESH MAULIK, AND ANDREW SNOWDEN

ABSTRACT. Given a finite field k of characteristic $p \geq 5$, we show that the Tate conjecture holds for K3 surfaces over $\overline{k}$ if and only if there are only finitely many K3 surfaces defined over each finite extension of k.

CONTENTS

1. Introduction 1
2. Tate implies finiteness 2
2.1. Discriminant bounds for the étale and crystalline lattices 2
2.2. Controlling the Néron–Severi lattice 5
2.3. Constructing low-degree ample line bundles 6
2.4. Finiteness of twisted forms 7
2.5. Proof of Main Theorem (1) 9
3. Finiteness implies Tate 9
3.1. Twisted sheaves 9
3.2. ℓ-adic B-fields 9
3.3. Twisted ℓ-adic Mukai lattices 11
3.4. Moduli spaces of twisted sheaves 13
3.5. Twisted partners of K3 surfaces over a finite field 15
3.6. Proof of Main Theorem (2) 19
References 19

1. INTRODUCTION

Given a class of algebraic varieties, it is reasonable to ask if there are only finitely many members defined over a given finite field. While this is clearly the case when the appropriate moduli functor is bounded, matters are often not so simple. For example, consider the case of abelian varieties of a given dimension g. There is no single moduli space parameterizing them; rather, for each integer $d \geq 1$ there is a moduli space parameterizing abelian varieties of dimension g with a polarization of degree d. It is nevertheless possible to show (see [Z, Theorem 4.1], [Mi, Corollary 13.13]) that there are only finitely many abelian varieties over a given finite field, up to isomorphism. Another natural class of varieties where this difficulty arises is the case of K3 surfaces. As with abelian varieties, there is not a single moduli space but rather a moduli space for each even integer $d \geq 2$, parametrizing K3 surfaces with a polarization of degree d.

In this paper, we consider the finiteness question for K3 surfaces over finite fields. Given a K3 surface X defined over a finite field k of characteristic p, the Tate conjecture predicts that the natural map

$$\mathrm{Pic}(X) \otimes \mathbf{Q}_\ell \to \mathrm{H}^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Q}_\ell(1))^{\mathrm{Gal}(\overline{k}/k)}$$

Date: November 15, 2011.

is surjective for $\ell \neq p$. It admits many alternate formulations; for example, it is equivalent to the statement that the Brauer group of X is finite. We say that X/k *satisfies the Tate conjecture over some extension* k'/k *(resp.* $\overline{k}$*)* if the Tate conjecture holds for the base change $X_{k'}$ (resp. for all base changes $X_{k'}$ with k'/k finite).

Our main result is that this conjecture is essentially equivalent to the finiteness of the set of K3 surfaces over k. Precisely:

Main Theorem. *Let k be a finite field of characteristic $p \geq 5$.*

1. *There are only finitely many isomorphism classes of K3 surfaces over k which satisfy the Tate conjecture over $\overline{k}$.*
2. *If there are only finitely many isomorphism classes of K3 surfaces over the quadratic extension k' of k then every K3 surface over k satisfies the Tate conjecture over k'.*

In particular, the Tate conjecture holds for all K3 surfaces over $\overline{k}$ if and only if there are only finitely many K3 surfaces defined over each finite extension of k.

As the Tate conjecture is known for K3 surfaces of finite height in characteristic at least 5 [NO], we obtain the following unconditional corollary:

Corollary. *There are only finitely many isomorphism classes of K3 surfaces of finite height defined over k.*

Our argument proceeds as follows. To obtain finiteness from Tate, it suffices to prove the existence of low-degree polarizations on K3 surfaces over k. In order to do this, we use the Tate conjecture in both ℓ-adic and crystalline cohomology to control the possibilities of the Néron-Severi lattice. For the other direction, we use the finiteness statement and the existence of infinitely many Brauer classes to create a K3 surface with infinitely many twisted Fourier-Mukai partners. Since this cannot happen in characteristic zero, we obtain a contradiction by proving a lifting result. This argument does not rely on [NO] (as it did in an earlier version of this paper).

Notation. Throughout, k denotes a finite field of characteristic p and cardinality $q = p^f$. We fix an algebraic closure $\overline{k}$ of k.

Acknowledgments. We would like to thank Jean-Louis Colliot-Thélène, Daniel Huybrechts, Abhinav Kumar, Matthias Schütt, Damiano Testa, Yuri Zarhin, and the referees for many helpful comments and discussions. M.L. is partially supported by NSF grant DMS-1021444, NSF CAREER grant DMS-1056129, and the Sloan Foundation. D.M. is partially supported by a Clay Research Fellowship.

2. TATE IMPLIES FINITENESS

2.1. **Discriminant bounds for the étale and crystalline lattices.** In this section, we produce bounds on the discriminants of certain lattices constructed from the étale and crystalline cohomologies of K3 surfaces over k. We begin by recalling some terminology. Let A be a principal ideal domain. By a *lattice* over A, we mean a finite free A module M together with a symmetric A-linear form $(,) : M \otimes_A M \to A$. We say that M is *non-degenerate* (resp. *unimodular*) if the map $M \to \mathrm{Hom}_A(M, A)$ provided by the pairing is injective (resp. bijective). The *discriminant* of a lattice M, denote $\mathrm{disc}(M)$, is the determinant of the matrix (e_i, e_j), where $\{e_i\}$ is a basis for M as an A-module; it is a well-defined element of $A/(A^\times)^2$. The lattice M is non-degenerate (resp. unimodular) if and only if its discriminant is non-zero (resp. a unit). Note that the valuation of $\mathrm{disc}(M)$ at a maximal ideal of A is well-defined.

We will need a simple lemma concerning discriminants:

Lemma 2.1.1. *Let A be a discrete valuation ring with uniformizer t. Let M be a lattice over A and let $M' \subset M$ be an A-submodule such that M/M' has length r as an A-module. Regard M' as a lattice by restricting the form from M. Then $\operatorname{disc}(M') = t^{2r} \operatorname{disc}(M)$ up to units of A.*

Proof. Let $e_1, \dots, e_n$ be a basis for M and let $f_1, \dots, f_n$ be a basis for M'. Let B be the matrix (e_i, e_j) and let B' be the matrix (f_i, f_j). Thus $\operatorname{disc}(M) = \det B$ and $\operatorname{disc}(M') = \det B'$. Let $C \in M_n(A)$ be the change of basis matrix, so that $f_i = Ce_i$. Then $\det(C) = t^r$ up to units of A. As $B' = C^t BC$, the result follows. ☐

The following general result on discriminant bounds will be used several times in what follows.

Proposition 2.1.2. *Fix a positive integer r and a non-negative even integer w. There exist constants C and C', depending only on r, w and q, with the following property.*

Let E be a finite unramified extension of $\mathbf{Q}_\ell$ with ring of integers $\mathcal{O}$. Let M be a lattice over $\mathcal{O}$ of rank r equipped with an endomorphism ϕ. Let v_0 be the ℓ-adic valuation of $\operatorname{disc}(M)$. Assume that the characteristic polynomial of ϕ belongs to $\mathbf{Z}[T]$, that all eigenvalues of ϕ on $M[1/\ell]$ are Weil q-integers of weight w and that $q^{w/2}$ is a semi-simple eigenvalue of ϕ on $M[1/\ell]$.

1. *If $\ell > C$ then the discriminant $M^{\phi=q^{w/2}}$ has ℓ-adic valuation at most v_0.*
2. *The discriminant of $M^{\phi=q^{w/2}}$ has ℓ-adic valuation at most $C' + v_0$.*

Proof. We first define the constants C and C'. Let $\mathcal{W}$ be the set of all Weil q-integers of weight w and degree at most r. It is easy to bound the coefficients of the minimal polynomial of an element of $\mathcal{W}$, and so one sees that $\mathcal{W}$ is a finite set. Let S denote the set of elements of $\mathbf{Z}[T]$ which are monic of degree r and whose roots belong to $\mathcal{W}$. Clearly, S is a finite set; enumerate its elements as $f_1(T), \dots, f_m(T)$. We can factor each $f_i(T)$ as $g_i(T)h_i(T)$, where $g_i(T)$ is a power of $T - q^{w/2}$ and $h_i(T)$ is an element of $\mathbf{Z}[T]$ which does not have $q^{w/2}$ as a root. For each i, pick rational polynomials $a_i(T)$ and $b_i(T)$ such that

$$a_i(T)g_i(T) + b_i(T)h_i(T) = 1.$$

Let Q be the least common multiple of the denominators of the coefficients of $a_i(T)$ and $b_i(T)$. Let s be the maximal integer such that ℓ^s divides Q, for some prime ℓ, and let ℓ_0 be the largest prime dividing Q. We claim that we can take $C = \ell_0$ and $C' = 2rs$.

We now prove these claims. Thus let M and ϕ be given, and put $N = M^{\phi=q^{w/2}}$. The characteristic polynomial of ϕ belongs to S, and is thus equal to $f_i(T)$ for some i. Put $M_1 = h_i(\phi)M$ and $M_2 = g_i(\phi)M$. One easily sees that $M_1 \oplus M_2$ is a finite index $\mathcal{O}$-submodule of M and that M_1 and M_2 are orthogonal. Furthermore, M_1 is contained in N, since $q^{w/2}$ is a semi-simple eigenvalue of ϕ, and has finite index.

Suppose that $\ell > C$. Then $a_i(T)$ and $b_i(T)$ belong to $\mathcal{O}[T]$ and so $M = M_1 \oplus M_2$. Thus $\operatorname{disc}(M) = \operatorname{disc}(M_1) \operatorname{disc}(M_2)$. It follows that $\operatorname{disc}(M_1)$ has ℓ-adic valuation at most v_0. As N and M_1 are saturated in M and $M_1 \subset N$, we have $N = M_1$, and so (1) follows.

Now suppose that ℓ is arbitrary. Then $\ell^s a_i(T)$ and $\ell^s b_i(T)$ belong to $\mathcal{O}[T]$. It follows that $M_1 \oplus M_2$ contains $\ell^s M$, and so $M/(M_1 \oplus M_2)$ has length at most rs as an $\mathcal{O}$-module. Lemma 2.1.1 shows that $\operatorname{disc}(M_1) \operatorname{disc}(M_2)$ divides $\ell^{2rs} \operatorname{disc}(M)$, and thus has ℓ-adic valuation as most $2rs + v_0 = C' + v_0$. The lemma also shows that $\operatorname{disc}(N)$ divides $\operatorname{disc}(M_1)$, which proves (2). ☐

Let X be a K3 surface over k. For a prime number $\ell \neq p$ put

$$M_\ell(X) = \mathrm{H}^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Z}_\ell), \qquad N_\ell(X) = M_\ell(X)^{\phi=q}.$$

Then $M_\ell(X)$ is a free $\mathbf{Z}_\ell$-module of rank 22, and the cup product gives it the structure of a unimodular lattice. The space $M_\ell(X)$ admits a natural $\mathbf{Z}_\ell$-linear automorphism ϕ, the geometric Frobenius element of $\operatorname{Gal}(\overline{k}/k)$. The map ϕ does not quite preserve the form, but satisfies

$(\phi x, \phi y) = q^2(x, y)$. It is known [D] that the action of ϕ on $M_\ell(X)$ is semi-simple. We give $N_\ell(X)$ the structure of a lattice by restricting the form from $M_\ell(X)$.

Proposition 2.1.3. *There exist constants $C_1 = C_1(k)$ and $C_2 = C_2(k)$ with the following properties. Let X be a K3 surface over k and let $\ell \neq p$ be a prime number. Then*

1. *For $\ell > C_1$, the discriminant of $N_\ell(X)$ has ℓ-adic valuation zero.*
2. *The discriminant of $N_\ell(X)$ has ℓ-adic valuation at most C_2.*

Proof. This follows immediately from Proposition 2.1.2 with $r = 22$ and $w = 2$, applied to $M = M_\ell(X)$. Note that $v_0 = 0$. ☐

We also need a version of the above result at p. Let $W = W(k)$ be the Witt ring of k. Put

$$M_p(X) = \mathrm{H}^2_{\mathrm{cris}}(X/W), \qquad N_p(X) = M_p(X)^{\phi_0 = p}.$$

Then $M_p(X)$ is a free W-module of rank 22, and the cup product gives it the structure of a unimodular lattice. The lattice $M_p(X)$ admits a natural semilinear automorphism ϕ_0, the crystalline Frobenius. The map $\phi = \phi_0^f$ is W-linear (where $q = p^f$). We have $(\phi_0 x, \phi_0 y) = p^2 \phi_0((x, y))$. (Note: the ϕ_0 on the right is the Frobenius on W.) Since ϕ_0 is only semi-linear, $N_p(X)$ is not a W-module, but a $\mathbf{Z}_p$-module. We give $N_p(X)$ the structure of a lattice via the form on $M_p(X)$.

We say that an eigenvalue α of a linear map is semi-simple if the α-eigenspace coincides with the α-generalized eigenspace. We now come to the main result at p:

Proposition 2.1.4. *There exists a constant $C_3 = C_3(k)$ with the following property. Let X be a K3 surface over k. Assume that q is a semi-simple eigenvalue of ϕ on $M_p(X)[1/p]$. Then the discriminant of $N_p(X)$ has p-adic valuation at most C_3.*

Proof. Let X be given, and put $N' = M_p(X)^{\phi = q}$, so that $N_p(X) = (N')^{\phi_0 = p}$. Proposition 2.1.2, with $r = 22$ and $w = 2$, bounds the p-adic valuation of $\mathrm{disc}(N')$ (as a lattice over W) in terms of k; in fact, the produced bound is the number C_2 from the previous proposition. The following lemma (which defines a constant C_4) now shows that the p-adic valuation of $\mathrm{disc}(N_p(X))$ is bounded by $C_3 = C_2 f + 44 C_4 f$. ☐

Lemma 2.1.5. *There exists a constant $C_4 = C_4(k)$ with the following property. Let N' be a lattice over W of rank n. Let ϕ_0 be a semi-linear endomorphism of N' satisfying $\phi_0^f = q$ and $(\phi_0 x, \phi_0 y) = p^2 \phi_0((x, y))$, and put $N = (N')^{\phi_0 = p}$. Then $v_p(\mathrm{disc}(N))$ is at most $f v_p(\mathrm{disc}(N')) + 2 C_4 n f$.*

Proof. Let $g(T) = T - p$ and let $h(T) = (T^f - q)/(T - p)$, two monic polynomials in $\mathbf{Z}_p[T]$. Pick polynomials $a(T)$ and $b(T)$ in $\mathbf{Q}_p[T]$ such that

$$a(T) g(T) + b(T) h(T) = 1.$$

Let r be such that $p^r a(T)$ and $p^r b(T)$ belong to $\mathbf{Z}_p[T]$. We claim that we can take $C_4 = r$.

Thus let N' with ϕ_0 be given. Let L' be N', regarded as a $\mathbf{Z}_p$-lattice with pairing $\langle , \rangle$ given by $\langle x, y \rangle = \mathrm{tr}_{W/\mathbf{Z}_p}(x, y)$. We have $\mathrm{rk}(L') = fn$ and $\mathrm{disc}(L') = \mathbf{N}(\mathrm{disc}(N'))$, where $\mathbf{N}$ is the norm from W to $\mathbf{Z}_p$. We regard ϕ_0 as a linear map of L'. As such, it is semi-simple (when p is inverted) with minimal polynomial $T^f - q$. The identity $\langle \phi_0 x, \phi_0 y \rangle = p^2 \langle x, y \rangle$ holds. Let L be the p eigenspace of ϕ_0, regarded as a sublattice of L'. Then L and N are the same subset of N', but the form on L is that on N scaled by f. It follows that $\mathrm{disc}(N)$ divides $\mathrm{disc}(L)$.

Put $L_1 = h(\phi_0) L'$ and $L_2 = g(\phi_0) L'$. Then L_1 and L_2 are orthogonal under $\langle , \rangle$ and $L_1 \oplus L_2$ has index at most p^{fnr} in L'. It follows from Lemma 2.1.1 that $\mathrm{disc}(L_1)$ divides $p^{2fnr} \mathrm{disc}(L')$. Since L_1 is contained in L, we find that $\mathrm{disc}(L)$ divides $p^{2fnr} \mathrm{disc}(L')$ as well. Finally, we see that $\mathrm{disc}(N)$ divides $p^{2fnr} \mathbf{N}(\mathrm{disc}(N'))$, and so $v_p(\mathrm{disc}(N))$ is bounded by $2fnr + f v_p(\mathrm{disc}\, N')$. ☐

Remark 2.1.6. As far as we are aware, it is not known if the action of ϕ on $M_p(X)[1/p]$ is semi-simple. See Corollary 7.5 of [O2] for a partial result. However, for K3 surfaces which satisfy the Tate conjecture, semi-simplicity is known, and can be easily be deduced from the aforementioned result of Ogus.

2.2. Controlling the Néron–Severi lattice.

We assume for the rest of §2 that $p \geq 5$. Let X be a K3 surface over k. Write $\mathrm{NS}(X)$ for the Néron–Severi group of X, which is a lattice (over $\mathbf{Z}$) under the intersection pairing. The main result of this section is the following:

Proposition 2.2.1. *There exists a finite set $\mathscr{L} = \mathscr{L}(k)$ of lattices over $\mathbf{Z}$ with the following property: if X is a K3 surface over k which satisfies the Tate conjecture (over k) then $\mathrm{NS}(X)$ is isomorphic to a member of $\mathscr{L}$.*

We handle the finite and infinite height cases separately. In the finite height case, the Tate conjecture is known. We will need integral refinements of it, both at ℓ and at p, which we now give.

Lemma 2.2.2. *Let X be a K3 surface over k which satisfies the Tate conjecture and let $\ell \neq p$ be a prime. Then the map*

$$c_1 : \mathrm{NS}(X) \otimes \mathbf{Z}_\ell \to N_\ell(X)(1)$$

is an isomorphism.

Proof. The Kummer sequence on $X_{\overline{k}}$ gives a short exact sequence

$$0 \to \mathrm{NS}(X_{\overline{k}}) \otimes \mathbf{Z}/\ell^n\mathbf{Z} \to \mathrm{H}^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Z}/\ell^n\mathbf{Z}(1)) \to \mathrm{Br}(X_{\overline{k}})[\ell^n] \to 0.$$

Taking the inverse limit over n, we obtain an exact sequence

$$0 \to \mathrm{NS}(X_{\overline{k}}) \otimes \mathbf{Z}_\ell \to \mathrm{H}^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Z}_\ell(1)) \to T_\ell(\mathrm{Br}(X_{\overline{k}})) \to 0,$$

where T_ℓ here denotes Tate module. The first map above is c_1. Since the ℓ-adic Tate module of any abelian group has no ℓ-torsion, we see that the image of $\mathrm{NS}(X_{\overline{k}}) \otimes \mathbf{Z}_\ell$ in $\mathrm{H}^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Z}_\ell(1))$ is saturated. As $\mathrm{NS}(X)$ is saturated in $\mathrm{NS}(X_{\overline{k}})$, it follows that the image of the map

$$c_1 : \mathrm{NS}(X) \otimes \mathbf{Z}_\ell \to N_\ell(X)(1)$$

is saturated. Since X satisfies the Tate conjecture, the above map is an isomorphism when ℓ is inverted. From saturatedness, it is therefore an isomorphism without ℓ inverted. ☐

Remark 2.2.3. The Chern class map appearing in the statement of the lemma may look odd, as $\mathrm{NS}(X)$ carries no Galois action but it appears as if $N_\ell(X)(1)$ does. However, one should regard $N_\ell(X)$ as carrying an action by the inverse of the cyclotomic character (as it is a $\phi = q$ eigenspace), and so the Tate twist cancels this action.

Lemma 2.2.4. *Let X be a K3 surface of finite height over k which satisfies the Tate conjecture. Then the map*

$$c_1 : \mathrm{NS}(X) \otimes \mathbf{Z}_p \to N_p(X)\{1\}$$

is an isomorphism.

Remark 2.2.5. By [NO], a finite height K3 surface satisfies the Tate conjecture, so we could omit that hypothesis from the lemma. However, we wish to make clear the independence of our argument from [NO].

Proof. Let $W = W(k)$ be the Witt ring and let E be the field of fractions of W. Since X satisfies the Tate conjecture, c_1 is an isomorphism after inverting p. To show that c_1 itself is an isomorphism, it is enough to show that the image of $\mathrm{NS}(X) \otimes W\{-1\}$ in $\mathrm{H}^2_{\mathrm{cris}}(X/W)$ is saturated. (We write $\{1\}$ for the crystalline version of Tate twist, so that $W\{-1\}$ is W with ϕ_0 acting by $1 \mapsto p$ and filtration concentrated in degree 1.) Let $\mathfrak{X}$ be a lift of X to W such that the restriction map $\mathrm{NS}(\mathfrak{X}) \to \mathrm{NS}(X)$ is an isomorphism; such a lift exists by [LM, Cor 4.2]. Of course, it suffices to show that the image of $\mathrm{NS}(\mathfrak{X}) \otimes W\{-1\}$ in $\mathrm{H}^2_{\mathrm{cris}}(X/W)$ is saturated.

We now need to make use of some integral p-adic Hodge theory, specifically, the theories of Fontaine–Laffaille and Fontaine–Messing. See [BM, §3] for a review of these theories. For a strongly divisible module M over W, we let $T^\vee(M)$ be the $\mathbf{Z}_p$-lattice $\mathrm{Hom}_{W,\mathrm{Fil},\phi}(M, A_{\mathrm{cris}})$ and we let $T(M)$ be the $\mathbf{Z}_p$-dual of $T^\vee(M)$. The functor T gives an equivalence between the category of strongly divisible lattices and the category of Galois stable lattices in crystalline representations of restricted weight (this follows from the discussion in [BM, §3]). From this we see that an inclusion $M \subset M'$ of strongly divisible lattices is saturated if and only if $T(M) \subset T(M')$ is saturated.

Now, the map $c_1 : \mathrm{NS}(X) \otimes W\{-1\} \to \mathrm{H}^2_{\mathrm{cris}}(\mathfrak{X}/W)$ is a map of strongly divisible modules. Thus we can check that the image is saturated after applying T. We have a commutative diagram

$$\begin{tikzcd}
\mathrm{NS}(\mathfrak{X}) \otimes \mathbf{Z}_p(-1) \arrow[r] & \mathrm{H}^2_{\text{ét}}(\mathfrak{X}_{\overline{E}}, \mathbf{Z}_p) \\
\mathrm{NS}(\mathfrak{X}) \otimes T(W\{-1\}) \arrow[r] \arrow[u] & T(\mathrm{H}^2_{\mathrm{cris}}(X/W)) \arrow[u]
\end{tikzcd}$$

The left arrow is just the isomorphism $T(W\{-1\}) = \mathbf{Z}_p(-1)$, while the right arrow is the Fontaine–Messing isomorphism (which exists since $p-1$ is greater than the degree of the cohomology group in consideration, namely 2). The top arrow is the usual étale Chern class while the bottom arrow is T applied to the crystalline Chern class map. The commutativity of the diagram is the compatibility of the p-adic comparison map with Chern classes (see [FM, §6.3]). It thus suffices to show that the image of the top map is saturated. Now, $\mathrm{NS}(\mathfrak{X}) = \mathrm{NS}(\mathfrak{X}_E)$, and $\mathrm{NS}(\mathfrak{X}_E)$ is saturated in $\mathrm{NS}(\mathfrak{X}_{\overline{E}})$. Thus it suffices to show that $\mathrm{NS}(\mathfrak{X}_{\overline{E}})(-1)$ is saturated in $\mathrm{H}^2_{\text{ét}}(\mathfrak{X}_{\overline{E}}, \mathbf{Q}_p)$. However, this is well-known (replace $\overline{E}$ by $\mathbf{C}$ and use the exponential sequence). □

We now return to the proof of the proposition.

Proof of Proposition 2.2.1. Let X be a finite height K3 surface over k. Let $\ell \neq p$ be a prime number. By Lemma 2.2.2, $\mathrm{NS}(X) \otimes \mathbf{Z}_\ell$ is isomorphic, as a lattice, to $N_\ell(X)(1)$. Since $N_\ell(X)(1)$ is isomorphic, as a lattice, to $N_\ell(X)$, we find that $\mathrm{disc}(\mathrm{NS}(X))$ and $\mathrm{disc}(N_\ell(X))$ have the same ℓ-adic valuations. Similarly, appealing to Lemma 2.2.4, we find that $\mathrm{disc}(\mathrm{NS}(X))$ and $\mathrm{disc}(N_p(X))$ have the same p-adic valuations. Applying Propositions 2.1.3 and 2.1.4, we find that $|\mathrm{disc}(\mathrm{NS}(X))|$ is at most $p^{C_3} \prod_{\ell \leq C_1} \ell^{C_2}$. As there are only finitely many isomorphism classes of lattices of a given rank and discriminant [Cas, Ch. 9, Thm. 1.1], it follows that there are only finitely many possibilities for $\mathrm{NS}(X)$ (up to isomorphism).

Now let X be an infinite height K3 surface over k which satisfies the Tate conjecture. By [O, §1.7] the discriminant of $\mathrm{NS}(X_{\overline{k}})$ is $-p^a$ where $1 \leq a \leq 20$ is even. The Frobenius element ϕ of $\mathrm{Gal}(\overline{k}/k)$ acts on $\mathrm{NS}(X_{\overline{k}})$ with finite order; it is therefore semi-simple and its eigenvalues are roots of unity, i.e., Weil q-integers of weight 0. Since $\mathrm{NS}(X) = \mathrm{NS}(X_{\overline{k}})^{\phi=1}$, Proposition 2.1.2 shows that the ℓ-adic valuation of $\mathrm{disc}(\mathrm{NS}(X))$ is bounded in terms of q for each ℓ, and 0 for ℓ sufficiently large; thus $\mathrm{disc}(\mathrm{NS}(X))$ is bounded in terms of q. Appealing again to the finiteness of lattices with given rank and discriminant, we find that there are only finitely many possibilities for $\mathrm{NS}(X)$ (up to isomorphism). □

2.3. **Constructing low-degree ample line bundles.** Let K be the extension of k of degree

$$6983776800 = 2^5 \cdot 3^3 \cdot 5^2 \cdot 7 \cdot 11 \cdot 13 \cdot 17 \cdot 19.$$

The field K is relevant for the following reason (explained below): if X is a K3 surface over k then any line bundle on $X_{\overline{k}}$ descends to one on X_K. The purpose of this section is to prove the following proposition:

Proposition 2.3.1. *There exists a constant $C_5 = C_5(k)$ with the following property: if X is a K3 surface over k which satisfies the Tate conjecture over K then X admits an ample line bundle of degree at most C_5 defined over K.*

We begin with a lemma.

Lemma 2.3.2. *Let X and X' be K3 surfaces over algebraically closed fields such that $\mathrm{NS}(X)$ and $\mathrm{NS}(X')$ are isomorphic as lattices. Then the set of degrees of ample line bundles on X and X' coincide.*

Proof. Put $N = \mathrm{NS}(X)$ and $N_{\mathbf{R}} = N \otimes \mathbf{R}$. Let Δ be the set of elements δ in N such that $(\delta, \delta) = -2$. For $\delta \in \Delta$, let $r_\delta : N_{\mathbf{R}} \to N_{\mathbf{R}}$ be the reflection given by $r_\delta(x) = x + (x, \delta)\delta$. Let Γ be the group generated by the r_δ, with $\delta \in \Delta$. Finally, let V be the set of elements x in $N_{\mathbf{R}}$ such that $(x, x) > 0$ and $(x, \delta) \neq 0$ for all $\delta \in \Delta$. Then V is an open subset of $N_{\mathbf{R}}$ and the group $\pm\Gamma$ acts transitively on the set of connected components of V [O, Prop. 1.10]. Furthermore, there exists a unique connected component V_0 of V such that an element of N is ample if and only if it lies in V_0 [O, p. 371].

Let N', etc., be as above but for X'. Choose an isomorphism of lattices $i : N \to N'$. Clearly, i induces a bijection $\Delta \to \Delta'$ and a homeomorphism $V \to V'$. Thus $i(V_0)$ is some connected component of V'. We can find an element γ of $\pm\Gamma'$ such that $\gamma(i(V_0)) = V'_0$. Thus, replacing i by γi, we can assume that $i(V_0) = V'_0$. It then follows that i induces a bijection between the set of ample elements in N and the set of ample elements in N'. Since i preserves degree, this proves the lemma. □

Corollary 2.3.3. *For every lattice L there is an integer $d(L)$ with the following property: if X is a K3 surface over an algebraically closed field such that $\mathrm{NS}(X)$ is isomorphic to L then X admits an ample line bundle of degree $d(L)$.*

We now return to the proof of Proposition 2.3.1.

Proof of Proposition 2.3.1. Let X be a K3 surface over k. The Frobenius element ϕ of the absolute Galois group of k acts on $\mathrm{NS}(X_{\overline{k}})$ as a finite order endomorphism. It is therefore semi-simple. Furthermore, since its characteristic polynomial has degree at most 22 and has at least one eigenvalue equal to 1, its remaining eigenvalues which are roots of unity of degree at most 21. At this point, we use the defining property of K. By construction, any integer m with $\varphi(m) \leq 21$ must divide $N = \deg[K : k]$. It follows that $\phi^N = 1$ holds. In particular, we see that $\mathrm{NS}(X_{\overline{k}}) = \mathrm{NS}(X_K)$.

Let $\mathscr{L} = \mathscr{L}(K)$ be the set of lattices provided by Proposition 2.2.1. Let C_5 be the maximum value of $d(L)$ for $L \in \mathscr{L}$, where $d(L)$ is as defined in Corollary 2.3.3. Let X be a K3 surface over k satisfying the Tate conjecture over K. Then $\mathrm{NS}(X_K)$, and thus $\mathrm{NS}(X_{\overline{k}})$, belongs to $\mathscr{L}$. It follows that $X_{\overline{k}}$, and thus X_K, admits an ample line bundle of degree at most C_5. This proves the proposition. □

2.4. **Finiteness of twisted forms.** Recall that a *twisted form* of a K3 surface X/k is a K3 surface over k which is isomorphic to X over $\overline{k}$. The purpose of this section is to establish the following result.

Proposition 2.4.1. *Let X be a K3 surface over the finite field k. Then X has only finitely many twisted forms, up to isomorphism.*

The set of isomorphism classes of twisted forms of X is in bijection with the non-abelian cohomology set $\mathrm{H}^1(\mathrm{Gal}(\overline{k}/k), \mathrm{Aut}_{\overline{k}}(X_{\overline{k}}))$, and so to prove the proposition it suffices to show finiteness of this set. We begin with two lemmas. In what follows, $\widehat{\mathbf{Z}}$ denotes the profinite completion of $\mathbf{Z}$ and ϕ a topological generator. Suppose $\widehat{\mathbf{Z}}$ acts continuously on a discrete group E. A 1-cocycle for this action is given by an element $x \in E$ such that $xx^{\phi} \cdots x^{\phi^{n-1}} = 1$ for all sufficiently divisible integers n. Moreover, cocycles represented by x and y are cohomologous if there exists an element $h \in E$ such that $x = h^{-1}yh^{\phi}$

Lemma 2.4.2. *Let G and G' be discrete groups on which $\widehat{\mathbf{Z}}$ acts continuously and let $f : G \to G'$ be a $\widehat{\mathbf{Z}}$-equivariant homomorphism whose kernel is finite and whose image has finite index. If $\mathrm{H}^1(\widehat{\mathbf{Z}}, G')$ is finite then $\mathrm{H}^1(\widehat{\mathbf{Z}}, G)$ is finite.*

Proof. Let $x_1, \ldots, x_n \in G'$ be cocycles representing the elements of $\mathrm{H}^1(\widehat{\mathbf{Z}}, G')$ and $y_1, \ldots, y_j \in G'$ representatives for the right cosets of $f(G)$ in G'. Given a cocycle $x \in G$, there is i and $h \in G'$ such that

$$h^{-1}f(x)h^{\phi} = x_i.$$

Choosing j and $z \in G$ such that $h = f(z)y_j$ yields

$$y_j^{-1}f(z)^{-1}f(x)f(z)^{\phi}y_j^{\phi} = x_i,$$

whence

$$f(z^{-1}xz^{\phi}) = y_j x_i (y_j^{\phi})^{-1}.$$

Since f has finite kernel, each element $y_j x_i (y_j^{\phi})^{-1}$ has finitely many preimages in G, and we see that the union of this finite set of finite sets contains cocycles representing all of $\mathrm{H}^1(\widehat{\mathbf{Z}}, G)$, as desired. ☐

Lemma 2.4.3. *Let G be a discrete group and let $\widehat{\mathbf{Z}} \to G$ be a continuous homomorphism; regard $\widehat{\mathbf{Z}}$ as acting on G by inner automorphisms via the homomorphism. Assume that G has only finitely many conjugacy classes of finite order elements. Then $H^1(\widehat{\mathbf{Z}}, G)$ is finite.*

Proof. Let g be the image of ϕ under the map $\widehat{\mathbf{Z}} \to G$. Continuity forces g to have finite order. Because $\widehat{\mathbf{Z}}$ acts by conjugation, the cocycle condition for an element $x \in G$ simply amounts to xg having finite order, and x and y are cohomologous if xg and yg are conjugate. We thus find that multiplication by g gives a bijection between $\mathrm{H}^1(\widehat{\mathbf{Z}}, G)$ and the set of conjugacy classes of finite order elements of G. This completes the proof. ☐

We now prove the proposition.

Proof of Proposition 2.4.1. Let $N = \mathrm{NS}(X_{\overline{k}})$ and let $N' \subset N_{\mathbf{R}}$ be the nef cone. Let G' be the group of automorphisms of the lattice N which map N' to itself, let $G = \mathrm{Aut}_{\overline{k}}(X_{\overline{k}})$, let $G^{\circ} = \mathrm{Aut}_k(X_{\overline{k}})$, and let $\Gamma = \mathrm{Aut}(\overline{k}/k) \cong \widehat{\mathbf{Z}}$. Since the natural action of G° on N preserves N', there are homomorphisms

$$\Gamma \to G^{\circ} \to G'.$$

In addition, conjugation by the image of ϕ in G° preserves G and gives the natural Frobenius action on the automorphism group. Thus, the natural map

$$f : G \to G'$$

is $\widehat{\mathbf{Z}}$-equivariant with respect to the natural action on G and the conjugation action on G'. The map f has finite kernel and its image has finite index by [LM, Thm 6.1]. Furthermore, G' has finitely many conjugacy classes of finite order elements (see [T, §6]). The above two lemmas thus imply that $\mathrm{H}^1(\Gamma, G)$ is finite, which completes the proof. ☐

2.5. **Proof of Main Theorem (1).** We now complete the proof of the first part of the main theorem. Let M_d be the stack over k of pairs (X, L) where X is a K3 surface and L is a polarization of degree d. It follows from Artin's representability theorem that M_d is Deligne–Mumford and locally of finite type over k; since the third power of any polarization is very ample [SD], the stack is of finite type. Let C_5 be the constant produced by Proposition 2.3.1. Consider the diagram

$$\begin{tikzcd}
\coprod_{d=1}^{C_5} M_d(K) \arrow[r, "\alpha"] & \{\text{isomorphism classes of K3's over } K\} \\
& \{\text{isomorphism classes of K3's over } k \text{ satisfying Tate over } K\} \arrow[u, "\beta"]
\end{tikzcd}$$

By the definition of C_5, any element in the image of β is also in the image of α. Since the domain of α is finite, it follows that the image of β is finite. Any two elements of a fiber of β are twisted forms of each other, and so the fibers of β are finite by Proposition 2.4.1. We thus find that the domain of β is finite, which completes the proof.

3. FINITENESS IMPLIES TATE

3.1. **Twisted sheaves.** We use the notions and notation from [L1], [L2], and the references therein. Recall the basic definition. Fix a $\boldsymbol{\mu}_r$-gerbe over an algebraic space $\mathscr{X} \to Z$.

Definition 3.1.1. A sheaf $\mathscr{F}$ of (left) $\mathscr{O}_{\mathscr{X}}$-modules is *$\lambda$-fold $\mathscr{X}$-twisted* if the natural left inertial $\boldsymbol{\mu}_r$-action $\boldsymbol{\mu}_r \times \mathscr{F} \to \mathscr{F}$ of a section ρ of $\boldsymbol{\mu}_r$ is scalar multiplication by the ρ^λ.

Notation 3.1.2. Given a $\boldsymbol{\mu}_n$-gerbe $\mathscr{X} \to Z$, write $\mathrm{D}^{\mathrm{tw}}(\mathscr{X})$ for the derived category of perfect complexes of $\mathscr{X}$-twisted sheaves and $\mathrm{D}^{-\mathrm{tw}}(\mathscr{X})$ for the derived category of perfect complexes of (-1)-fold $\mathscr{X}$-twisted sheaves.

3.2. **ℓ-adic B-fields.** Let Z be a separated scheme of finite type over the field k. The following is an "ℓ-adification" of a notion familiar from mathematical physics. For the most part, this rephrases well-known results in a form that aligns them with the literature on twisted Mukai lattices, to be developed ℓ-adically in the next section.

Definition 3.2.1. An *ℓ-adic B-field on Z* is an element

$$B \in \mathrm{H}^2_{\text{ét}}(Z, \mathbf{Q}_\ell(1)).$$

We can write any B-field as α/ℓ^n with

$$\alpha \in \mathrm{H}^2_{\text{ét}}(Z, \mathbf{Z}_\ell(1))$$

a primitive element. *When we write B in this form we will always assume (unless noted otherwise) that α is primitive.*

Definition 3.2.2. Given a B-field α/ℓ^n on Z, the *Brauer class associated to B* is the image of α under the map

$$\mathrm{H}^2_{\text{ét}}(Z, \mathbf{Z}_\ell(1)) \to \mathrm{H}^2_{\text{ét}}(Z, \boldsymbol{\mu}_{\ell^n}) \to \mathrm{Br}(Z)[\ell^n].$$

Notation 3.2.3. Given $\alpha \in \mathrm{H}^2_{\text{ét}}(Z, \mathbf{Z}_\ell(1))$ we will write α_n for the image in $\mathrm{H}^2_{\text{ét}}(Z, \boldsymbol{\mu}_{\ell^n})$. We will use brackets to indicate the map $\mathrm{H}^2_{\text{ét}}(Z, \boldsymbol{\mu}_{\ell^n}) \to \mathrm{Br}(X)$.

Thus, the Brauer class associated to the B-field α/ℓ^n is written $[\alpha_n]$.

Lemma 3.2.4. *Given $\alpha \in \mathrm{H}^2_{\text{ét}}(Z, \mathbf{Z}_\ell(1))$ and positive integers n, n', we have that*

$$\ell^n[\alpha_{n+n'}] = [\alpha_{n'}] \in \mathrm{Br}(Z).$$

Proof. Consider the commutative diagram of Kummer sequences

$$\begin{tikzcd}
1 \arrow[r] & \boldsymbol{\mu}_{\ell^{n+n'}} \arrow[r] \arrow[d, "\ell^n"'] & \mathbf{G}_m \arrow[r, "\ell^{n+n'}"] \arrow[d, "\ell^n"'] & \mathbf{G}_m \arrow[r] \arrow[d, "\mathrm{id}"] & 1 \\
1 \arrow[r] & \boldsymbol{\mu}_{\ell^{n'}} \arrow[r] & \mathbf{G}_m \arrow[r, "\ell^{n'}"] & \mathbf{G}_m \arrow[r] & 1.
\end{tikzcd}$$

The induced map

$$\ell^n : \mathrm{H}^2_{\text{ét}}(Z, \boldsymbol{\mu}_{\ell^{n+n'}}) \to \mathrm{H}^2_{\text{ét}}(Z, \boldsymbol{\mu}_{\ell^{n'}})$$

is identified with the reduction map

$$\mathrm{H}^2_{\text{ét}}(Z, \mathbf{Z}_\ell(1) \otimes \mathbf{Z}/\ell^{n+n'}\mathbf{Z}) \to \mathrm{H}^2_{\text{ét}}(Z, \mathbf{Z}_\ell(1) \otimes \mathbf{Z}/\ell^{n'}\mathbf{Z}),$$

so it sends $\alpha_{n+n'}$ to $\alpha_{n'}$. On the other hand, ℓ^n acts by multiplication by ℓ^n on $\mathrm{H}^2_{\text{ét}}(Z, \mathbf{G}_m)$. The result follows from the resulting diagram of cohomology groups. ☐

Lemma 3.2.5. *The Brauer classes associated to ℓ-adic B-fields on X form a subgroup*

$$\mathrm{Br}^B_\ell(Z) \subset \mathrm{Br}(Z)(\ell)$$

of the ℓ-primary part of the Brauer group of Z.

Proof. Let α/ℓ^n and β/ℓ^m be B-fields with Brauer classes $[\alpha_n]$ and $[\beta_m]$. By Lemma 3.2.4 we have that $[\alpha_n] = \ell^m[\alpha_{n+m}]$ and $[\beta_m] = \ell^n[\beta_{n+m}]$. Let $\gamma = \ell^m\alpha + \ell^n\beta$. We have that

$$[\gamma_{n+m}] = \ell^m[\alpha_{n+m}] + \ell^n[\beta_{n+m}] = [\alpha_n] + [\beta_m],$$

as desired. ☐

Given a smooth projective geometrically connected algebraic surface X over k, the intersection pairing defines a map

$$\mathrm{H}^2_{\text{ét}}(X, \mathbf{Z}_\ell(1)) \times \mathrm{H}^2_{\text{ét}}(X, \mathbf{Z}_\ell(1)) \to \mathrm{H}^4_{\text{ét}}(X_{\overline{k}}, \mathbf{Z}_\ell(2)) = \mathbf{Z}_\ell.$$

There is a cycle class map

$$\mathrm{Pic}(X) \otimes \mathbf{Z}_\ell \to \mathrm{H}^2_{\text{ét}}(X, \mathbf{Z}_\ell(1)).$$

Write $P(X, \mathbf{Z}_\ell)$ for its image, which is a $\mathbf{Z}_\ell$-sublattice.

Definition 3.2.6. The *ℓ-adic transcendental lattice of X* is

$$T(X, \mathbf{Z}_\ell) := P(X, \mathbf{Z}_\ell)^\perp \subset \mathrm{H}^2_{\text{ét}}(X, \mathbf{Z}_\ell(1))$$

Lemma 3.2.7. *The map $\beta : \alpha \otimes (1/\ell^n) \mapsto [\alpha_n]$ defines an isomorphism*

$$T(X, \mathbf{Z}_\ell) \otimes \mathbf{Q}_\ell/\mathbf{Z}_\ell \xrightarrow{\sim} \mathrm{Br}^B_\ell(X).$$

Proof. The map β extends to a map

$$\beta : \mathrm{H}^2_{\text{ét}}(X, \mathbf{Z}_\ell(1)) \otimes \mathbf{Q}_\ell \to \mathrm{Br}_\ell(X)$$

by sending α/ℓ^n to the element $[\alpha_n]$ as above. One easily checks that this map is well-defined, and it is surjective by the definition of $\mathrm{Br}_\ell(X)$. Moreover, by construction

$$P(X, \mathbf{Z}_\ell) \otimes \mathbf{Q}_\ell + \mathrm{H}^2_{\text{ét}}(X, \mathbf{Z}_\ell(1))$$

lies in the kernel of β. Since

$$\mathrm{H}^2_{\text{ét}}(X, \mathbf{Z}_\ell(1)) \otimes \mathbf{Q}_\ell = P(X, \mathbf{Z}_\ell) \otimes \mathbf{Q}_\ell \oplus T(X, \mathbf{Z}_\ell) \otimes \mathbf{Q}_\ell,$$

we see that the map defined in the statement of the Lemma is surjective. It remains to show that it is injective.

Suppose α/ℓ^n maps to 0 in $\mathrm{Br}(X)$. We have that $[\alpha_n] = 0$, so that

$$\alpha_n \in \mathrm{Pic}(X)/\ell^n \,\mathrm{Pic}(X) \subset \mathrm{H}^2_{\text{ét}}(X, \boldsymbol{\mu}_{\ell^n}).$$

Taking the cohomology of the exact sequence

$$0 \to \mathbf{Z}_\ell(1) \to \mathbf{Z}_\ell(1) \to \boldsymbol{\mu}_{\ell^n} \to 0,$$

we see that

$$\alpha \in \ell^n \, \mathrm{H}^2_{\text{ét}}(X, \mathbf{Z}_\ell(1)),$$

so that

$$\alpha/\ell^n = 0 \in T(X, \mathbf{Z}_\ell) \otimes \mathbf{Q}_\ell/\mathbf{Z}_\ell,$$

as desired. □

Proposition 3.2.8. *If X is a smooth projective surface over the finite field k of characteristic p then the following are equivalent.*

(1) $\mathrm{Br}(X)$ *is infinite*
(2) $\mathrm{Br}^B_\ell(X) \neq 0$ *for all* ℓ
(3) $\mathrm{Br}^B_\ell(X) \neq 0$ *for some* ℓ
(4) $T(X, \mathbf{Z}_\ell) \neq 0$

Proof. By [LLR], $\mathrm{Br}(X)$ is infinite if and only if $\mathrm{Br}(X)(\ell)$ is infinite for one ℓ if and only if $\mathrm{Br}(X)(\ell)$ is infinite for all ℓ. To prove the Proposition it suffices to prove that $\mathrm{Br}^B_\ell(X) \neq 0$ if $\mathrm{Br}(X)(\ell)$ is infinite. So suppose $\alpha_n \in \mathrm{Br}(X)(\ell^n)$ is a sequence of classes. Choose lifts

$$\widetilde{\alpha}_n \in \mathrm{H}^2_{\text{ét}}(X, \boldsymbol{\mu}_{\ell^n})$$

for each n. Let $\beta_0 = 0 \in \mathrm{H}^2_{\text{ét}}(X, \boldsymbol{\mu}_{\ell^0})$. Assume we have constructed $\beta_1, \ldots, \beta_m$, $\beta_i \in \mathrm{H}^2_{\text{ét}}(X, \boldsymbol{\mu}_{\ell^i})$ such that $\ell\beta_{i+1} = \beta_i$ and $\ell^j \widetilde{\alpha}_{m+j} = \beta_m$ for all $j > 0$. The group $\mathrm{H}^2_{\text{ét}}(X, \boldsymbol{\mu}_{\ell^{m+1}})$ is finite, so there is one element β_{m+1} that is a multiple of infinitely many $\widetilde{\alpha}_n$ such that $\ell\beta_{m+1} = \beta_m$. Replacing the sequence of $\widetilde{\alpha}_n$ with the subsequence mapping to β_{m+1}, we see that we can proceed by induction, yielding an element

$$\beta \in \mathrm{H}^2_{\text{ét}}(X, \mathbf{Z}_\ell(1))$$

giving infinitely many distinct elements of $\mathrm{Br}^B_\ell(X)$. □

3.3. Twisted ℓ-adic Mukai lattices.

Fix a K3 surface over the field k and an element $\alpha \in T(X, \mathbf{Z}_\ell)$. Fix a B-field α/r with $r = \ell^n$ for some n. Note that because X is simply connected it follows from the Leray spectral sequence that

$$\mathrm{H}^4(X, \mathbf{Z}_\ell(2)) = \mathrm{H}^4(X_{\overline{k}}, \mathbf{Z}_\ell(2)) = \mathbf{Z}_\ell.$$

Definition 3.3.1. The ℓ-adic Mukai lattice of X is the free $\mathbf{Z}_\ell$-module

$$\mathrm{H}_{\text{ét}}(X, \mathbf{Z}_\ell) := \mathrm{H}^0_{\text{ét}}(X, \mathbf{Z}_\ell) \oplus \mathrm{H}^2_{\text{ét}}(X, \mathbf{Z}_\ell(1)) \oplus \mathrm{H}^4_{\text{ét}}(X, \mathbf{Z}_\ell(2))$$

with the intersection pairing

$$(a, b, c) \cdot (a', b', c') = bb' - ac' - a'c \in \mathrm{H}^4_{\text{ét}}(X, \mathbf{Z}_\ell(2)) = \mathbf{Z}_\ell.$$

The algebraic part of the cohomology gives a sublattice

$$\mathrm{CH}(X, \mathbf{Z}_\ell) = \mathbf{Z}_\ell \oplus P(X, \mathbf{Z}_\ell) \oplus \mathbf{Z}_\ell.$$

It is easy to see that

$$\mathrm{CH}(X, \mathbf{Z}_\ell)^\perp = T(X, \mathbf{Z}_\ell),$$

as sublattices of $\mathrm{H}_{\text{ét}}(X, \mathbf{Z}_\ell)$. We will write $\mathrm{H}_{\text{ét}}(X, \mathbf{Q}_\ell)$ (resp. $\mathrm{CH}(X, \mathbf{Q}_\ell)$, resp. $T(X, \mathbf{Q}_\ell)$) for $\mathrm{H}_{\text{ét}}(X, \mathbf{Z}_\ell) \otimes_{\mathbf{Z}_\ell} \mathbf{Q}_\ell$ (resp. $\mathrm{CH}(X, \mathbf{Z}_\ell) \otimes \mathbf{Q}_\ell$), resp. $T(X, \mathbf{Z}_\ell) \otimes \mathbf{Q}_\ell$). The lattice $\mathrm{CH}(X, \mathbf{Z}_\ell)$ has an integral structure:

$$\mathrm{CH}(X, \mathbf{Z}) = \mathbf{Z} \oplus \mathrm{Pic}(X) \oplus \mathbf{Z}$$

such that $\mathrm{CH}(X, \mathbf{Z}) \otimes \mathbf{Z}_\ell = \mathrm{CH}(X, \mathbf{Z}_\ell)$. We will also write $\mathrm{CH}(X, \mathbf{Q})$ for $\mathrm{CH}(X, \mathbf{Z}) \otimes \mathbf{Q}$.

Following [Y], we consider the map

$$T_{-\alpha/r} : \mathrm{H}_{\text{ét}}(X, \mathbf{Z}_\ell) \to \mathrm{H}_{\text{ét}}(X, \mathbf{Q}_\ell)$$

that sends x to the cup product $x \cup e^{-\alpha/r}$.

Definition 3.3.2. The *α/r-twisted Chow lattice of X* is

$$\mathrm{CH}^{\alpha/r}(X, \mathbf{Z}_\ell) := (T_{-\alpha/r})^{-1}(\mathrm{CH}(X, \mathbf{Q}_\ell)).$$

There is also an integral structure on the twisted Chow lattice.

Lemma 3.3.3. *Suppose α is primitive. Let*

$$\mathrm{CH}^{\alpha/r}(X, \mathbf{Z}) = \{(ar, D + a\alpha, c) \mid a, c \in \mathbf{Z}, D \in \mathrm{Pic}(X)\} \subset T^{-1}_{-\alpha/r}(\mathrm{CH}(X, \mathbf{Q})) \subset \mathrm{CH}^{\alpha/r}(X, \mathbf{Z}_\ell).$$

The natural map

$$\mathrm{CH}^{\alpha/r}(X, \mathbf{Z}) \otimes \mathbf{Z}_\ell \to \mathrm{CH}^{\alpha/r}(X, \mathbf{Z}_\ell)$$

is an isomorphism.

Proof. A basis for $\mathrm{CH}^{\alpha/r}(X, \mathbf{Z})$ is given by $(r, \alpha, 0)$, the vectors $(0, b, 0)$ where b ranges over a basis for $\mathrm{Pic}(X)$, and $(0, 0, 1)$. Since $\mathrm{Pic}(X)$ is a lattice, we see that these vectors all remain linearly independent over $\mathbf{Z}_\ell$ (inside $\mathrm{H}_{\text{ét}}(X, \mathbf{Z}_\ell)$), so that the displayed scalar extension map is injective. To see surjectivity, suppose $(x, y, z) \in \mathrm{H}_{\text{ét}}(Z, \mathbf{Z}_\ell)$ is an element such that $e^{-\alpha/r} \cup (x, y, z) \in \mathrm{CH}(X, \mathbf{Q}_\ell)$. Computing the components of the cup product shows that (x, y, z) must satisfy the conditions: $x, y \in \mathbf{Z}_\ell$ (via the natural identifications $\mathrm{H}^0(X, \mathbf{Z}_\ell) = \mathbf{Z}_\ell$ and $\mathrm{H}^4(X, \mathbf{Z}_\ell(2)) = \mathbf{Z}_\ell$) and $-\alpha x/r + y \in \mathrm{Pic}(X) \otimes \mathbf{Q}_\ell$. Since α is primitive and orthogonal to $\mathrm{Pic}(X)$ and $y \in \mathrm{H}^2(X, \mathbf{Z}_\ell)$, we see that we must have $x = ar$ for some a and then $-a\alpha + y \in \mathrm{Pic}(X) \otimes \mathbf{Z}_\ell$, so $y = a\alpha + D$ for some $D \in \mathrm{Pic}(X, \mathbf{Z}_\ell)$. This shows that the displayed map is surjective. □

The (integral) twisted Chow lattice is a natural recipient of Chern classes for twisted sheaves. Let $\pi : \mathscr{X} \to X$ be a $\boldsymbol{\mu}_r$-gerbe representing the class $-[\alpha_n]$ associated to the B-field $-\alpha/r$, $r = \ell^n$. Suppose $\mathscr{P}$ is a perfect complex of $\mathscr{X}$-twisted sheaves of positive rank.

Definition 3.3.4. The *Chern character of $\mathscr{P}$* is the unique element

$$\mathrm{ch}_{\mathscr{X}}(\mathscr{P}) \in \mathrm{CH}(X, \mathbf{Q})$$

such that

$$\mathrm{rk}\,\mathrm{ch}_{\mathscr{X}}(\mathscr{P}) = \mathrm{rk}\,\mathscr{P}$$

and

$$\mathrm{ch}_{\mathscr{X}}(\mathscr{P})^r = \mathrm{ch}\left(\mathbf{R}\pi_*(\mathscr{P}^{\overset{\mathbf{L}}{\otimes} r})\right) \in \mathrm{CH}(X, \mathbf{Q}).$$

Remark 3.3.5. A less *ad hoc* approach is to define rational Chern classes using a splitting principle, etc. This can be done and yields the same result (and works for complexes of rank 0). In particular, the Chern character defined here satisfies the Riemann-Roch theorem in the following sense: given two perfect complexes $\mathscr{P}$ and $\mathscr{Q}$ of $\mathscr{X}$-twisted sheaves, we can compute $\chi(\mathscr{H}om(\mathscr{P}, \mathscr{Q}))$ as $\deg(\mathrm{ch}(\mathscr{P}^\vee) \cdot \mathrm{ch}(\mathscr{Q}) \cdot \mathrm{Td}_X)$.

Definition 3.3.6. Given a perfect complex $\mathscr{P}$ of $\mathscr{X}$-twisted sheaves as above, we define the *twisted Mukai vector of $\mathscr{P}$* to be

$$v^{\alpha/r}(\mathscr{P}) := e^{\alpha/r}\,\mathrm{ch}_{\mathscr{X}}(\mathscr{P})\sqrt{\mathrm{Td}_X}.$$

Lemma 3.3.7. *Given perfect complexes $\mathscr{P}$ and $\mathscr{Q}$ of $\mathscr{X}$-twisted sheaves as above, the following hold.*

(1) *the element $v^{\alpha/r}(\mathscr{P})$ lies in the integral subring* $\mathrm{CH}^{\alpha/r}(X, \mathbf{Z})$

(2) *we have that*

$$\chi(\mathscr{P},\mathscr{Q}) = -v^{\alpha/r}(\mathscr{P})\cdot v^{\alpha/r}(\mathscr{Q}).$$

Proof. To see the integrality, we may assume that the base field is algebraically closed. Since the calculation of $v^{\alpha/r}$ factors through K-theory, it is enough to show the analogous result for a class in the integral K-theory of twisted sheaves. Note that $\det\mathscr{P}$ is naturally the pullback of an invertible sheaf on X, since $\mathscr{P}$ has rank divisible by ℓ^n (α being primitive by assumption). We can thus work with the universal deformation of X over which a chosen ample divisor H and $\det\mathscr{P}$ remain algebraic, so that the class of $\mathscr{P}$ in K-theory deforms. In particular, we may prove the result after deforming to characteristic 0. Since the integrality is invariant under algebraically closed base field extension, we may assume that the base field is $\mathbf{C}$. Lemma 3.1, Lemma 3.3, and Remark 3.2 of [Y] then imply that $v^{\alpha/r}(\mathscr{P})$ has precisely the form described in Lemma 3.3.3.

To see the second statement, note that

$$\begin{aligned}-v^{\alpha/r}(\mathscr{P})\cdot v^{\alpha/r}(\mathscr{Q}) &= \deg\left(\mathrm{ch}_{\mathscr{X}}(\mathscr{P}^\vee \overset{\mathbf{L}}{\otimes}\mathscr{Q})\,\mathrm{Td}_X\right)\\ &= \deg\left(\mathrm{ch}(\mathbf{R}\pi_*(\mathbf{R}\mathscr{H}om(\mathscr{P},\mathscr{Q}))\,\mathrm{Td}_X\right)\\ &= \chi(\mathbf{R}\pi_*\mathbf{R}\mathscr{H}om(\mathscr{P},\mathscr{Q}))\\ &= \chi(\mathscr{P},\mathscr{Q})\end{aligned}$$

□

3.4. **Moduli spaces of twisted sheaves.** We recall the essential details of the moduli theory of twisted sheaves on K3 surfaces that we will need, building on the foundational work in [Mu]. The reader can consult [Y] for a development of a more general theory using a stack-free formulation of the notion of twisted sheaf. In particular, we only consider the case in which the rank equals the order of the Brauer class in order to simplify the exposition; [Y] contains the general theory (over $\mathbf{C}$).

Fix a K3 surface X over k, α/ℓ^n a primitive ℓ-adic B-field, and $v=(r,D,s)\in\mathrm{CH}^{\alpha/\ell^n}(X,\mathbf{Z})$ a Mukai vector.

Definition 3.4.1. The *stack of simple $\mathscr{X}$-twisted sheaves with Mukai vector* v is the stack $\mathscr{M}_{\mathscr{X}}(v)$ whose objects over a k-scheme T are pairs $(\mathscr{F},\phi)$, where $\mathscr{F}$ is a T-flat quasi-coherent $\mathscr{X}_T$-twisted sheaf of finite presentation and $\phi:\det\mathscr{F}\xrightarrow{\sim}\mathscr{O}(D)$ is an isomorphism of invertible sheaves on $\mathscr{X}$, such that for every geometric point $t\to T$, the fiber sheaf $\mathscr{F}_t$ has Mukai vector v and endomorphism ring $k(t)$.

Using Artin's Representability Theorem, one can show that $\mathscr{M}_{\mathscr{X}}(v)$ is an Artin stack locally of finite type over k. This is proven in Section 2.3.1 of [L1].

Proposition 3.4.2. *Given an ℓ-adic B-field α/ℓ^n and a primitive Mukai vector*

$$v\in\mathrm{CH}^{\alpha/\ell^n}(X,\mathbf{Z})$$

such that $\mathrm{rk}\,v=\ell^n$ *and* $v^2=0$*, we have that*

1. *the stack $\mathscr{M}_{\mathscr{X}}(v)$ of simple $\mathscr{X}$-twisted sheaves with Mukai vector v is a $\boldsymbol{\mu}_{\ell^n}$-gerbe over a K3 surface $M_{\mathscr{X}}(v)$;*
2. *the universal sheaf $\mathscr{E}$ on $\mathscr{X}\times\mathscr{M}_{\mathscr{X}}(v)$ defines an equivalence of derived categories*

$$\mathrm{D}^{\mathrm{tw}}(\mathscr{X})\xrightarrow{\sim}\mathrm{D}^{-\mathrm{tw}}(\mathscr{M}_{\mathscr{X}}(v));$$

3. *if there is a vector $u\in\mathrm{CH}^{\alpha/\ell^n}(X,\mathbf{Z})$ such that $(u\cdot v,\ell)=1$ then the Brauer class of the gerbe*

$$\mathscr{M}_{\mathscr{X}}(v)\to M_{\mathscr{X}}(v)$$

is trivial. In particular, there is an equivalence of derived categories

$$\mathrm{D}^{-\mathrm{tw}}(\mathscr{M}_{\mathscr{X}}(v)) \xrightarrow{\sim} \mathrm{D}(M_{\mathscr{X}}(v)).$$

Proof. It suffices to prove the first statement after extending scalars to $\overline{k}$. First note that since the sheaves in question have rank ℓ^n and the order of the Brauer class is ℓ^n, every object is automatically stable with respect to any polarization: any subsheaf has rank either 0 or ℓ^n, so stability conditions become vacuous. (See, e.g., Sections 2.2.5 and 3.2.4 of [L1] for further details.)

Let L be the second component of v. Tensoring the sheaves parametrized by $\mathscr{M}_{\mathscr{X}}(v)$ by an invertible sheaf M results in an isomorphic moduli problem parametrizing sheaves with determinant $L \otimes M^{\otimes \ell^n}$. Thus, taking $M \in \operatorname{Pic}(X) \setminus p \operatorname{Pic}(X)$ to be sufficiently ample we see that we can assume that L is ample and not contained in $p \operatorname{Pic}(X)$.

By Deligne's theorem [D] and the smooth and proper base change theorems in étale cohomology, there is a pointed connected smooth $W(k)$-scheme (M, m), a projective relative K3 surface $\mathcal{X} \to M$, a class $\widetilde{v} = (\ell^n, \mathscr{L}, c) \in \mathrm{CH}(\mathcal{X})$, a class $\widetilde{\alpha} \in \mathrm{H}^2(\mathcal{X}, \mathbf{Z}_\ell(1))$, and an isomorphism between the fiber of the triple $(\mathcal{X}, \widetilde{v}, \widetilde{\alpha})$ over m and the triple (X, v, α). Moreover, the geometric generic fiber of $\mathcal{X} \to M$ has Picard group generated by the restriction of $\mathscr{L}$. We choose a Henselian discrete valuation ring R with algebraically closed residue field dominating M and centered at m and pull the family back to R. This results in a triple $(\mathcal{X}_R, \widetilde{v}_R, \widetilde{\alpha}_R)$ with closed fiber (X, v, α) (up to algebraically closed base field extension) and geometric generic fiber having Picard number 1 and characteristic 0, on which v remains primitive. Write H for the relative polarization from $\mathscr{L}$ and v for $\widetilde{v}_R$.

Consider the relative moduli space $\mathscr{M}(v) \to \operatorname{Spec} R$ parametrizing H-stable $\mathscr{X}$-twisted sheaves with Mukai vector v. Since any simple twisted sheaf with unobstructed determinant is unobstructed, we know that $\mathscr{M}(v)$ is a $\boldsymbol{\mu}_{\ell^n}$-gerbe over a smooth algebraic space $\mathcal{M} \to \operatorname{Spec} R$. By Theorem 3.16 of [Y], the geometric generic fiber of $\mathcal{M}$ over R is a K3 surface. By Lemma 2.3.3.2 of [L1], the algebraic space $\mathcal{M}$ is proper over R. It follows that $\mathcal{M} \to R$ is a relative K3 surface, as claimed. In particular, the special fiber is a K3 surface.

By the foundational results on Fourier-Mukai transforms (see, for example, [Căl]), the second statement is equivalent to the adjunction maps

$$\mathscr{O}_{\Delta_{\mathscr{X}}} \to \mathbf{R}(\mathrm{pr}_{13})_* \left(\text{Ł} \, \mathrm{pr}_{12}^* \, \mathscr{E} \overset{\mathbf{L}}{\otimes} \text{Ł} \, \mathrm{pr}_{23}^* \, \mathscr{E}^\vee \right) \tag{3.4.2.1}$$

and

$$\mathbf{R}(\mathrm{pr}_{13})_* \left(\text{Ł} \, \mathrm{pr}_{12}^* \, \mathscr{E} \overset{\mathbf{L}}{\otimes} \text{Ł} \, \mathrm{pr}_{23}^* \, \mathscr{E}^\vee \right) \to \mathscr{O}_{\Delta_{\mathscr{X}}} \tag{3.4.2.2}$$

being quasi-isomorphisms. (See Proposition 3.3 of [LO] for a proof without any assumptions on the base field.) It thus suffices to prove the result after base change to $\overline{k}$. In this case, the proof proceeds precisely as in section 5.2 (proof of Theorem 1.2) of [Căl]: one verifies the classical criterion of Bridgeland for the Fourier-Mukai functor to give an equivalence of derived categories.

We now return to the situation in which our base field is the original finite field k. To prove the last statement, it suffices to prove the following lemma.

Lemma 3.4.3. *Given a Mukai vector*

$$u \in \mathrm{CH}^{\alpha/\ell^n}(X, \mathbf{Z}),$$

there is a perfect complex $\mathscr{P}$ of $\mathscr{X}$-twisted sheaves such that

$$v^{\alpha/\ell^n}(\mathscr{P}) = u.$$

Let us accept the lemma for a moment and see why this implies the result. Write $\mathscr{E}$ for the universal sheaf on

$$\mathscr{X} \times \mathscr{M}_{\mathscr{X}}(v),$$

and let p and q denote the first and second projections of that product, respectively. The sheaf $\mathscr{E}$ is simultaneously $\mathscr{X}$- and $\mathscr{M}_{\mathscr{X}}(v)$-twisted. Given a complex $\mathscr{P}$ as in Lemma 3.4.3, consider the perfect complex of $\mathscr{M}_{\mathscr{X}}(v)$-twisted sheaves

$$\mathscr{Q} := \mathbf{R}q_*(\text{Ł}p^*\mathscr{P}^\vee \overset{\mathbf{L}}{\otimes} \mathscr{E}).$$

The rank of this complex over a geometric point m of $\mathscr{M}$ is calculated by

$$\chi(\mathscr{P}, \mathscr{E}_m) = -v^{\alpha/\ell^n}(\mathscr{P}) \cdot v = -u \cdot v,$$

which is relatively prime to ℓ. By standard results, we have that the Brauer class of $\mathscr{M}_{\mathscr{X}}(v)$ satisfies

$$[\mathscr{M}_{\mathscr{X}}(v)] \in \mathrm{Br}(M_{\mathscr{X}}(v))[u \cdot v].$$

On the other hand, $\mathscr{M}_{\mathscr{X}}(v) \to M_{\mathscr{X}}(v)$ is a $\boldsymbol{\mu}_{\ell^n}$-gerbe, which implies that

$$[\mathscr{M}_{\mathscr{X}}(v)] \in \mathrm{Br}(M_{\mathscr{X}}(v))[\ell^n].$$

Combining the two statements yields the result.

It remains to prove Lemma 3.4.3.

Proof of Lemma 3.4.3. We know that

$$u = (ra, D + a\alpha, c),$$

so we seek a perfect complex of rank ra, determinant D, and appropriate second Chern class. By Theorem 4.3.1.1 of [L2], there is a locally free $\mathscr{X}$-twisted sheaf V of rank r. Moreover, since any curve over a finite field has trivial Brauer group the restriction $\mathscr{X} \times_X C$ has trivial Brauer class, hence supports twisted sheaves L_C of rank 1 by Lemma 3.1.1.8 of [L2]. Since $\det L_C \cong \mathscr{O}(C)$, we see that we can get any determinant by adding (in the derived category) a sum of shifts of invertible sheaves supported on curves. Finally, Wedderburn's theorem yields invertible twisted sheaves supported at any closed point p of X, and they have twisted Mukai vector $(0, 0, d)$, where d is the degree of p. By the Lang-Weil estimates [LW] X has a 0-cycle of degree 1, so there is a sum of (shifted) twisted skyscraper sheaves giving the desired second Chern class. Since any bounded complex on $\mathscr{X}$ is perfect ($\mathscr{X}$ being regular), the lemma is proven. ☐

This completes the proof of Proposition 3.4.2. ☐

Remark 3.4.4. We note that Theorem 4.3.1.1 of [L2] is far from trivial; it relies on de Jong's period-index theorem and a careful analysis of the asymptotic geometry of moduli spaces of twisted sheaves on surfaces (in particular, their asymptotic irreducibility). In this sense, the last part of Proposition 3.4.2, giving a "numerical" criterion linking a twisted derived category to an untwisted one, is the deepest. As we will see below, this equivalence is the glue that holds this direction of the argument together.

3.5. **Twisted partners of K3 surfaces over a finite field.** Fix a K3 surface X over the finite field k of characteristic $p \geq 5$. Let k'/k be the quadratic extension of k.

Lemma 3.5.1. *Let V be a non-degenerate quadratic space over a field F and let ϕ be a semi-simple element of the orthogonal group of V, all of whose eigenvalues belong to F. Then*

$$\mathrm{disc}(V^{\phi=1} \oplus V^{\phi=-1}) = (-1)^{n/2}\,\mathrm{disc}(V),$$

where $n = \dim(V) - \dim(V^{\phi=1} \oplus V^{\phi=-1})$.

Proof. Write V_λ for the λ eigenspace of ϕ. Suppose $\lambda \neq \pm 1$. Since ϕ preserves the form, elements of V_λ pair to zero with elements of V_μ unless $\mu = \lambda^{-1}$. Note that $\lambda^{-1} \neq \lambda$. The discriminant of $V_\lambda \oplus V_{\lambda^{-1}}$ is $(-1)^{d/2}$, where $d = \dim(V_\lambda)$, as is easily seen by choosing a basis for V_λ, taking the dual basis for $V_{\lambda^{-1}}$ and computing the matrix of pairings. Now, list the eigenvalues of ϕ, other than ± 1, as $\lambda_1, \lambda_1^{-1}, \ldots, \lambda_k, \lambda_k^{-1}$. Then V decomposes as a direct sum of V_1, V_{-1} and the $V_{\lambda_i} \oplus V_{\lambda_i^{-1}}$. We thus find that $\operatorname{disc}(V) = \operatorname{disc}(V_1 \oplus V_{-1})(-1)^{n/2}$, as was to be shown. □

Lemma 3.5.2. *Suppose that* $\operatorname{Br}(X_{k'})$ *is infinite, and let* d *be a rational number. Then there are infinitely many primes* ℓ *for which* $T(X_{k'}, \mathbf{Z}_\ell)$ *contains an element* γ *with* $\gamma^2 = d$.

Proof. Let $\phi' = \phi^2$ be the Frobenius over k'. If ℓ does not divide the discriminant of $\operatorname{Pic}(X_{k'})$ then the discriminant of $P(X_{k'}, \mathbf{Z}_\ell)$ is an ℓ-adic unit, and we have an orthogonal decomposition

$$H^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Z}_\ell(1))^{\phi'=1} = P(X_{k'}, \mathbf{Z}_\ell) \oplus T(X_{k'}, \mathbf{Z}_\ell). \tag{3.5.2.3}$$

By Proposition 2.1.3, the discriminant of the lattice $H^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Z}_\ell(1))^{\phi'=1}$ is an ℓ-adic unit for $\ell \gg 0$. We thus find that the discriminant of $T(X_{k'}, \mathbf{Z}_\ell)$ is an ℓ-adic unit for $\ell \gg 0$. Therefore, if $T(X_{k'}, \mathbf{Z}_\ell)$ has rank at least two then any element of $\mathbf{Z}_\ell$ is of the form γ^2 for some $\gamma \in T(X_{k'}, \mathbf{Z}_\ell)$. (This follows from [S, §1.7, Prop. 3] and Hensel's lemma.)

We must now handle the case where the transcendental lattice has rank one. Let E be the number field generated by the eigenvalues of ϕ and the square roots of -1, d and the discriminant of $\operatorname{Pic}(X_{k'})$. Let $\ell \gg 0$ be a large prime which splits completely in E. Recall from [D] that the action of ϕ on $H^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Q}_\ell(1))$ is semi-simple. We have

$$H^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Q}_\ell(1))^{\phi'=1} = H^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Q}_\ell(1))^{\phi=1} \oplus H^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Q}_\ell(1))^{\phi=-1}$$

Lemma 3.5.1 thus shows that the discriminant of $H^2_{\text{ét}}(X_{\overline{k}}, \mathbf{Q}_\ell(1))^{\phi'=1}$ is ± 1, and therefore a square in $\mathbf{Q}_\ell$; the same is true when we use $\mathbf{Z}_\ell$ coefficients. Since the discriminant of $\operatorname{Pic}(X_{k'})$ is also a square in $\mathbf{Q}_\ell$, the equation (3.5.2.3) shows that the discriminant of $T(X_{k'}, \mathbf{Z}_\ell)$ is a square in $\mathbf{Q}_\ell$. As in the previous paragraph, we know this discriminant is also a unit. Since the transcendental lattice has rank 1, we thus find that it has an element α such that $\alpha^2 = 1$. Since d is a square in $\mathbf{Q}_\ell$, we can find a multiple γ of α with $\gamma^2 = d$. □

Now assume that X is a K3 surface over the finite field k such that $\operatorname{Br}(X)$ is infinite. Fix a primitive ample divisor $D \in \operatorname{Pic}(X)$ and choose an odd prime ℓ relatively prime to $\operatorname{disc}\operatorname{Pic}(X)$ and D^2 and an element $\gamma \in T(X_{k'}, \mathbf{Z}_\ell)$ such that $\gamma^2 = -D^2$. (This is possible by the above lemma.)

Definition 3.5.3. Given a relative K3 surface $\mathcal{Y} \to S$, an invertible sheaf $\mathscr{L} \in \operatorname{Pic}(\mathcal{Y})$, and positive integers r and s, let

$$\mathbf{Sh}_{\mathcal{Y}/S}(w) \to S$$

be the stack of simple locally free sheaves with Mukai vector $w = (r, \mathscr{L}, s)$ on each fiber.

BE CLear about determinant

Lemma 3.5.4. *If* w *is primitive and* $w^2 = 0$ *then* $\mathbf{Sh}_{\mathcal{Y}/S}(w)$ *is a* $\boldsymbol{\mu}_r$*-gerbe over a smooth algebraic space* $\operatorname{Sh}_{\mathcal{Y}/S}(w)$ *of relative dimension* 2 *over* S *with non-empty geometric fibers.*

YO. clarify how Y is used, etc. Non-emptiness of the geometric fibers is proven in Section 3 of [Y]. The smoothness follows from the fact that that obstruction theory of a locally free sheaf V with determinant $\mathscr{L}$ on a fiber $\mathcal{Y}_s$ is given by the kernel of the trace map

$$\operatorname{Ext}^2(V, V) \to \mathrm{H}^2_{\text{ét}}(\mathcal{Y}_s, \mathscr{O}),$$

which vanishes when V is simple. The relative dimension is

$$\dim \operatorname{Ext}^1(V,V) = w^2 + 2 = 2,$$

as desired. □

When $S = \operatorname{Spec} L$ and the base field is understood, we will write simply $\mathbf{Sh}_{\mathcal{Y}}(w)$ and $\mathrm{Sh}_{\mathcal{Y}}(w)$.

Proposition 3.5.5. *In the above situation, there is an infinite sequence of pairs* (γ_n, M_n) *such that*

1. $\gamma_n \in \mathrm{H}^2_{\text{ét}}(X_{k'}, \boldsymbol{\mu}_{\ell^n})$ *with* $[\gamma_n]$ *of exact order* ℓ^n *in* $\mathrm{Br}(X_{\overline{k}})$*;*
2. M_n *is a K3 surface over* k' *such that* $\rho(M_n) = \rho(X_{k'})$ *and*
$$\operatorname{rk} T(M_n, \mathbf{Z}_\ell) = \operatorname{rk} T(X_{k'}, \mathbf{Z}_\ell);$$
3. *given a gerbe* $\mathscr{X}_n \to X_{k'}$ *representing* γ_n*, we have that* M_n *is a fine moduli space of stable* $\mathscr{X}_n$*-twisted sheaves with determinant* D*;*
4. *for each* n*, there is a (classical) Mukai vector* $w_n \in \mathrm{CH}(M_n)$ *such that*
   - (a) *a tautological sheaf* $\mathscr{E}$ *on* $\mathscr{X}_n \times M_n$ *induces an open immersion*
   $$\mathscr{X}_n \hookrightarrow \mathbf{Sh}_{M_n}(w_n);$$
   - (b) *given a complete dvr* R *with residue field* k' *and a relative K3 surface* $\mathcal{M}$ *over* R *with* $\mathcal{M} \otimes_R k' = M_n$ *such that* $\operatorname{Pic}(\mathcal{M}) = \operatorname{Pic}(M_n)$*, there is a relative K3 surface* $\mathcal{X} \to \operatorname{Spec} R$ *such that* $\mathcal{X} \otimes k' = X$*, a* $\boldsymbol{\mu}_{\ell^n}$*-gerbe* $\mathfrak{X} \to \mathcal{X}$ *such that* $\mathfrak{X} \otimes_R K = \mathscr{X}_n$*, and a locally free* $\mathfrak{X} \times \mathcal{M}$*-twisted sheaf* $\mathfrak{E}$ *such that* $\mathfrak{E} \otimes_R k' = \mathscr{E}$*. For any inclusion* $R \hookrightarrow \mathbf{C}$*, the base changed family* $(\mathfrak{X}_{\mathbf{C}}, \mathcal{M}_{\mathbf{C}}, \mathfrak{E}_{\mathbf{C}})$ *yields a twisted Fourier-Mukai partnership*
   $$(\mathcal{X}_{\mathbf{C}}, [\mathfrak{X}_{\mathbf{C}}]) \sim \mathcal{M}_{\mathbf{C}},$$
   where $[\mathfrak{X}_{\mathbf{C}}] \in \mathrm{Br}(\mathcal{X}_{\mathbf{C}})$ *is a Brauer class of exact order* ℓ^n*.*

Proof. Consider the Mukai vector

$$v_n := (\ell^n, \gamma + D, 0) \in \mathrm{CH}^{\gamma/\ell^n}(X_{k'}, \mathbf{Z}).$$

We have that

$$v_n^2 = \gamma^2 + D^2 = 2q^2 + D^2 = 0$$

and

$$v_n \cdot (\ell^n, \gamma, 0) = \gamma^2 = 2q^2 = -D^2 \not\equiv 0 \mod \ell.$$

It is easy to see that γ is primitive. Consider the diagram

$$\begin{tikzcd}
0 \arrow[r] & \operatorname{Pic}(X_{k'}) \otimes \mathbf{Z}_\ell \arrow[r] \arrow[d] & \mathrm{H}^2(X_{k'}, \mathbf{Z}_\ell(1)) \arrow[r] \arrow[d] & T_\ell \operatorname{Br}(X_{k'}) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \operatorname{Pic}(X_{\overline{k}}) \otimes \mathbf{Z}_\ell \arrow[r] & \mathrm{H}^2(X_{\overline{k}}, \mathbf{Z}_\ell(1)) \arrow[r] & T_\ell \operatorname{Br}(X_{\overline{k}}) \arrow[r] & 0,
\end{tikzcd}$$

where T_ℓ denotes "ℓ-adic Tate module".

We claim that the left square is Cartesian. Indeed, it follows from the Hochschild-Serre spectral sequence that the middle vertical arrow identifies the top group with the Frobenius-invariants in the bottom group. Similarly, since the Brauer obstruction to the existence of invertible sheaves on proper varieties over finite fields is trivial, the left vertical arrow is also the group invariants. But the invariants in a submodule are just the intersection with the ambient invariants, showing that the square is Cartesian. It follows that for each n, the image of γ_n in $\mathrm{Br}(X_{\overline{k}})$ has order exactly ℓ^n, as claimed.

Let $\mathscr{X}_n$ be a gerbe representing $-\gamma_n$ (see Notation 3.2.3 for the definition of γ_n in terms of γ). Applying Proposition 3.4.2, we have that $\mathscr{M}_{\mathscr{X}_n}(v_n)$ is a $\boldsymbol{\mu}_{\ell^n}$-gerbe over a K3 surface M_n whose Brauer class is killed by

$$v_n \cdot (\ell^n, \gamma, 0) = -D^2,$$

hence is trivial. This yields an equivalence

$$\mathrm{D}^{\mathrm{tw}}(\mathscr{X}_n) \xrightarrow{\sim} \mathrm{D}(M_n).$$

Finally, the universal sheaf $\mathscr{E}$ on $\mathscr{X}_n \times \mathscr{M}_{\mathscr{X}_n}(v_n)$ induces an isometry

$$\mathrm{H}_{\text{ét}}(X_{k'}, \mathbf{Q}_\ell) \xrightarrow{\sim} \mathrm{H}_{\text{ét}}(M_n, \mathbf{Q}_\ell)$$

that restricts to an isometry

$$\mathrm{CH}(X_{k'}, \mathbf{Q}_\ell) \xrightarrow{\sim} \mathrm{CH}(M_n, \mathbf{Q}_\ell)$$

(see Section 4.1.1 of [Y] for the proof, written using Yoshioka's notation), establishing the first three statements of the Proposition.

Let $\mathscr{E}$ be a tautological sheaf on $\mathscr{X}_n \times M_n$. The determinant of $\mathscr{E}$ is naturally identified with the pullback of an invertible sheaf

$$L_{X_n} \boxtimes L_{M_n} \in \mathrm{Pic}(X_n \times M_n),$$

and each geometric fiber $\mathscr{E}_x$ has a second Chern class $c \in \mathbf{Z}$. Letting

$$w_n = (\ell^n, L_{M_n}, c),$$

the sheaf $\mathscr{E}$ gives a morphism of $\boldsymbol{\mu}_{\ell^n}$-gerbes

$$\mathscr{X}_n \to \mathbf{Sh}_{M_n}(w_n).$$

Since $\mathscr{E}$ is a Fourier-Mukai kernel, this morphism is an étale monomorphism (see e.g. Section 4 of [LO]), so it is an open immersion.

Now fix a lift $\mathcal{M}$ of M_n over R over which all of $\mathrm{Pic}(M_n)$ lifts (by assumption). Since $\mathrm{Pic}(M_n)$ lifts, so does w_n and $\mathbf{Sh}_{\mathcal{M}/R}(w_n)$ is a $\boldsymbol{\mu}_{\ell^n}$-gerbe over a smooth algebraic space $\mathrm{Sh}_{\mathcal{M}/R}(w_n)$ over R. Write $\mathscr{V}$ for the universal sheaf on $\mathbf{Sh}_{\mathcal{M}/R}(w_n) \times_R \mathcal{M}$. We can write

$$\det \mathscr{V} = \mathscr{U} \boxtimes L_{M_n}$$

with

$$\mathscr{U} \in \mathrm{Pic}(\mathrm{Sh}_{\mathcal{M}/R}(w_n)),$$

and by assumption the pullback of $\mathscr{U}$ along the map

$$X_{k'} \to \mathrm{Sh}_{M_n}(w_n) \hookrightarrow \mathrm{Sh}_{\mathcal{M}/R}(w_n)$$

is isomorphic to D, hence is ample (by the assumption on D). Since $\mathscr{X}_n$ is open in $\mathbf{Sh}_{M_n}(w_n)$, the induced open formal substack

$$\mathscr{Z} \subset \widehat{\mathbf{Sh}}_{\mathcal{M}/R}(w_n) \to \mathrm{Spf}\, R$$

is a $\boldsymbol{\mu}_{\ell^n}$-gerbe over a formal deformation $\mathfrak{Z}$ of $X_{k'}$ over $\mathrm{Spf}\, R$. The determinant of the universal formal sheaf gives a formal lift of D over $\mathfrak{Z}$, hence a polarization. It follows that $\mathscr{Z} \to \mathfrak{Z}$ is algebraizable, giving a $\boldsymbol{\mu}_{\ell^n}$-gerbe over a relative K3 surface

$$\mathscr{G} \to \mathcal{X} \to \mathrm{Spec}\, R$$

and an open immersion

$$\mathscr{G} \hookrightarrow \mathbf{Sh}_{\mathcal{M}/R}(w_n)$$

extending

$$\mathscr{X}_n \hookrightarrow \mathbf{Sh}_{M_n}(w_n).$$

Restricting the universal sheaf gives the desired deformation $\mathfrak{E}$ of $\mathscr{E}$. Nakayama's lemma shows that $\mathfrak{E}$ is the kernel of a relative Fourier-Mukai equivalence (i.e., the adjunction maps

(3.4.2.1) and (3.4.2.2) are quasi-isomorphisms bceause their derived restrictions to the fiber are quasi-isomorphisms). Foundational details are contained in Section 3 of [LO], and a similar deformation argument is contained in Section 6 of [LO]. □

3.6. **Proof of Main Theorem (2).** Fix a finite field k of characteristic p, and let k' be the quadratic extension of k. We now prove the second part of the main theorem from the introduction.

Theorem 3.6.1. *Suppose* $p \geq 5$. *If there are only finitely many K3 surfaces over* k' *then for any K3 surface* X *over* k*, the Brauer group* $X_{k'}$ *is finite, i.e., the Tate conjecture holds for* $X_{k'}$.

Proof. Suppose that X is a K3 surface over k such that $\mathrm{Br}(X_{k'})$ is infinite. We know by Proposition 3.5.5 that there is an infinite sequence of Brauer classes $\gamma_n \in \mathrm{Br}(X_{k'})$ such that

1. γ_n has exact order ℓ^n over $\overline{k}$, and
2. the twisted K3 surface $(X_{k'}, \gamma_n)$ is Fourier-Mukai equivalent to a K3 surface M_n over k'.

If there are only finitely many K3 surfaces over k', there is a subsequence n_i such that all of the M_{n_i} are the same surface, say M. Since M has infinite Brauer group (by Proposition 3.5.5(2)), it is not Shioda-supersingular. It follows from Proposition 3.1 and Proposition 4.1 of [LM] that there is a lift $\mathcal{M}$ of M over a complete dvr R with residue field finite over k' such that $\mathrm{Pic}(\mathcal{M}) = \mathrm{Pic}(M)$. Applying Proposition 3.5.5(4)(b) we end up with a complex K3 surface $\mathcal{M}_{\mathbf{C}}$ with a sequence of twisted derived partners (X_i, η_i) with $\eta_i \in \mathrm{Br}(X_i)$ of exact order ℓ^{n_i} for a strictly increasing sequence n_i. In particular, $\mathcal{M}_{\mathbf{C}}$ has infinitely many twisted partners. But this is a contradiction by Corollary 4.6 of [HS]. □

REFERENCES

[BM] C. Breuil and W. Messing, *Torsion étale and crystalline cohomologies*, Cohomologies p-adiques et applications arithmétiques, II., Astérisque No. 279 (2002), 81–124.

[Căl] A. Căldăraru, *Nonfine moduli spaces of sheaves on K3 surfaces*, Int. Math. Res. Not. 2002, no. 20, 1027–1056.

[Cas] J. W. S. Cassels, *Rational quadratic forms*, London Mathematical Society Monographs, 13, Academic Press, Inc., 1978.

[D] P. Deligne, *La conjecture de Weil pour les surfaces K3*, Invent. Math. **15** (1972), 206–226.

[FM] J.-M. Fontaine and W. Messing, *p-adic periods and p-adic étale cohomology*, Current trends in arithmetical algebraic geometry (Arcata, Calif., 1985), 179–207, Contemp. Math., **67**, Amer. Math. Soc., Providence, RI, 1987.

[HS] D. Huybrechts and P. Stellari, *Equivalences of twisted K3 surfaces*, Math. Ann. 332 (2005), no. 4, 901–936.

[LW] S. Lang and A. Weil, *Number of points of varieties in nite elds*, Amer. J. Math., 76:819–827, 1954

[L1] M. Lieblich, *Moduli of twisted sheaves*, Duke Math. J. **138** (2007), no. 1, 23–118.

[L2] M. Lieblich, *Twisted sheaves and the period-index problem*, Compos. Math. **144** (2008), no. 1, 1–31.

[LM] M. Lieblich and D. Maulik, *A note on the cone conjecture for K3 surfaces in positive characteristic*, preprint, 2011, arXiv:1102.3377.

[LO] M. Lieblich and M. Olsson, *Derived equivalence of K3 surfaces in positive characteristic*, in preparation.

[LLR] Q. Liu, D. Lorenzini, and M. Raynaud, *On the Brauer group of a surface*, Invent. Math. **159** (2005), no. 3, 673–676.

[Mi] J. S. Milne, *Abelian varieties*, v2.00, 2008, available at `www.jmilne.org/math/`.

[Mu] S. Mukai, *On the moduli space of bundles on K3 surfaces. I.*, Vector bundles on algebraic varieties (Bombay, 1984), 341–413, Tata Inst. Fund. Res. Stud. Math., 11, Tata Inst. Fund. Res., Bombay, 1987.

[NO] N. Nygaard and A. Ogus, *Tate's conjecture for K3 surfaces of finite height*, Ann. of Math. (2) **122** (1985), no. 3, 461–507

[O] A. Ogus, *A Crystalline Torelli Theorem for Supersingular K3 Surfaces*, in "Arithmetic and Geometry," Progress in Mathematics **36**, Birkhäuser 1983.

[O2] A. Ogus, *F-isocrystals and de Rham cohomology. II. Convergent isocrystals.*, Duke Math. J. **51** (1984), no. 4, 765–850.

[S] J.-P. Serre, *A course in arithmetic*, Graduate Texts in Mathematics, No. 7. Springer–Verlag, 1973.
[SD] B. Saint-Donat, *Projective models of K3 surfaces*, Amer. J. Math. **96** (1974), 602–639.
[T] B. Totaro, *Algebraic surfaces and hyperbolic geometry*, preprint, 2010, arXiv:1008.3825.
[Y] K. Yoshioka, *Moduli spaces of twisted sheaves on a projective variety*, Moduli spaces and arithmetic geometry, 1–30, Adv. Stud. Pure Math., 45, Math. Soc. Japan, Tokyo, 2006.
[Z] Ju. .G. Zarhin, *Endomorphisms of abelian varieties and points of finite order in characteristic P* (Russian), Mat. Zametki **21** (1977), no. 6, 737–744.

E-mail address: `lieblich@math.washington.edu`

E-mail address: `dmaulik@math.mit.edu`

E-mail address: `asnowden@math.mit.edu`